


Path integral molecular dynamics

David Manolopoulos
Department of Chemistry, University of Oxford



$$\Delta q = \Lambda(T) / \sqrt{8\pi}$$


$$\Lambda(T) = h / \sqrt{2\pi m k T}$$

$$\Delta p = \sqrt{m k T}$$

$$\Rightarrow \Delta p \Delta q = \frac{\hbar}{2}$$

Mariapfarr Workshop 2019, Lecture I

I. The classical isomorphism^{1,2}



$$Q = \frac{1}{(2\pi\hbar)^n} \int d\mathbf{p} \int d\mathbf{q} e^{-\beta_n H_n(\mathbf{p}, \mathbf{q})}$$

$$H_n(\mathbf{p}, \mathbf{q}) = \sum_{j=1}^n \left[\frac{p_j^2}{2m} + \frac{1}{2} m \omega_n^2 (q_j - q_{j+1})^2 + V(q_j) \right]; \quad \beta_n = \beta/n; \quad \omega_n = 1/(\beta_n \hbar).$$

Background:

In standard (basis set) quantum mechanics

$$\langle \nu | \nu' \rangle = \delta_{\nu\nu'} \quad \hat{1} = \sum_{\nu} |\nu\rangle \langle \nu| \quad \text{tr} [\hat{A}] = \sum_{\nu} \langle \nu | \hat{A} | \nu \rangle .$$

In the position and momentum representations³

$$\langle q | p \rangle = \sqrt{\frac{1}{2\pi\hbar}} e^{+ipq/\hbar} \quad \langle p | q \rangle = \langle q | p \rangle^* = \sqrt{\frac{1}{2\pi\hbar}} e^{-ipq/\hbar}$$

the analogous results are

$$\begin{aligned} \langle q | q' \rangle &= \delta(q - q') & \langle p | p' \rangle &= \delta(p - p') \\ \hat{1} &= \int dq |q\rangle \langle q| & \hat{1} &= \int dp |p\rangle \langle p| \\ \text{tr} [\hat{A}] &= \int dq \langle q | \hat{A} | q \rangle & \text{tr} [\hat{A}] &= \int dp \langle p | \hat{A} | p \rangle . \end{aligned}$$

Proof of the isomorphism:

$$Q = \text{tr} \left[e^{-\beta \hat{H}} \right] = \text{tr} \left[\left(e^{-\beta_n \hat{H}} \right)^n \right] \text{ where } \beta_n = \beta/n.$$

So

$$Q = \int dq_1 \dots \int dq_n \langle q_1 | e^{-\beta_n \hat{H}} | q_2 \rangle \dots \langle q_n | e^{-\beta_n \hat{H}} | q_1 \rangle,$$

with

$$\begin{aligned} \langle q | e^{-\beta_n \hat{H}} | q' \rangle &\simeq \langle q | e^{-\beta_n \hat{V}/2} e^{-\beta_n \hat{T}} e^{-\beta_n \hat{V}/2} | q' \rangle \\ &= \frac{1}{2\pi\hbar} \int dp e^{-\beta_n p^2/2m + ip(q-q')/\hbar - \beta_n [V(q)/2 + V(q')/2]} \\ &= \frac{1}{2\pi\hbar} \left(\frac{2\pi m}{\beta_n} \right)^{1/2} e^{-\beta_n [m\omega_n^2 (q-q')^2/2 + V(q)/2 + V(q')/2]} \\ &= \frac{1}{2\pi\hbar} \int dp e^{-\beta_n [p^2/2m + m\omega_n^2 (q-q')^2/2 + V(q)/2 + V(q')/2]}, \end{aligned}$$

gives Q with an error of $O(n^{-2})$.

2. Path integral molecular dynamics⁴

PIMD uses the ring polymer trajectories

$$\dot{\mathbf{q}} = + \frac{\partial H_n(\mathbf{p}, \mathbf{q})}{\partial \mathbf{p}} \quad \dot{\mathbf{p}} = - \frac{\partial H_n(\mathbf{p}, \mathbf{q})}{\partial \mathbf{q}}$$

as a sampling tool to calculate *exact* values of static equilibrium properties such as

$$\langle A \rangle = \frac{1}{Q} \text{tr} \left[e^{-\beta \hat{H}} \hat{A} \right].$$

Average potential energy

$$\begin{aligned}\text{tr} \left[e^{-\beta \hat{H}} \hat{V} \right] &= \text{tr} \left[\left(e^{-\beta_n \hat{H}} \right)^{j-1} \hat{V} \left(e^{-\beta_n \hat{H}} \right)^{n+1-j} \right] \\ &= \int dq_1 \dots \int dq_n \langle q_1 | e^{-\beta_n \hat{H}} | q_2 \rangle \dots \langle q_j | \hat{V} e^{-\beta_n \hat{H}} | q_{j+1} \rangle \dots \langle q_n | e^{-\beta_n \hat{H}} | q_1 \rangle \\ &= \int dq_1 \dots \int dq_n \langle q_1 | e^{-\beta_n \hat{H}} | q_2 \rangle \dots \langle q_n | e^{-\beta_n \hat{H}} | q_1 \rangle V(q_j) \\ &= \frac{1}{(2\pi\hbar)^n} \int d\mathbf{p} \int d\mathbf{q} e^{-\beta_n H_n(\mathbf{p}, \mathbf{q})} V(q_j)\end{aligned}$$


for any bead $j = 1, 2, \dots, n$, and we can improve the statistics by averaging over the beads:

$$\langle V \rangle = \frac{1}{(2\pi\hbar)^n Q} \int d\mathbf{p} \int d\mathbf{q} e^{-\beta_n H_n(\mathbf{p}, \mathbf{q})} \mathcal{V}(\mathbf{q}) \equiv \langle \mathcal{V}(\mathbf{q}) \rangle,$$

where the potential energy estimator is

$$\mathcal{V}(\mathbf{q}) = \frac{1}{n} \sum_{j=1}^n V(q_j).$$

Average kinetic energy



By the same argument, the kinetic energy estimator can be constructed from

$$\langle q_j | e^{-\beta_n \hat{H}/2} \hat{T} e^{-\beta_n \hat{H}/2} | q_{j+1} \rangle \simeq \left[\frac{1}{2\beta_n} - \frac{1}{2} m \omega_n^2 (q_j - q_{j+1})^2 \right] \langle q_j | e^{-\beta_n \hat{H}} | q_{j+1} \rangle$$

for any bead j , and averaging over the beads to improve the statistics gives

$$\langle T \rangle = \frac{1}{(2\pi\hbar)^n Q} \int d\mathbf{p} \int d\mathbf{q} e^{-\beta_n H_n(\mathbf{p}, \mathbf{q})} \mathcal{T}(\mathbf{q}) \equiv \langle \mathcal{T}(\mathbf{q}) \rangle$$

where the kinetic energy estimator is

$$\mathcal{T}(\mathbf{q}) = \frac{1}{2\beta_n} - \frac{1}{2n} \sum_{j=1}^n m \omega_n^2 (q_j - q_{j+1})^2.$$

Thermodynamic energy estimator

An alternative approach is to note that the average value of the energy $E = T + V$ is given by statistical mechanics as

$$\langle E \rangle = - \left(\frac{\partial \ln Q}{\partial \beta} \right)_V = - \frac{1}{Q} \left(\frac{\partial Q}{\partial \beta} \right)_V$$

which with

$$Q = \frac{1}{(2\pi\hbar)^n} \int d\mathbf{p} \int d\mathbf{q} e^{-\beta_n H_n(\mathbf{p}, \mathbf{q})}$$

and

$$H_n(\mathbf{p}, \mathbf{q}) = \sum_{j=1}^n \left[\frac{p_j^2}{2m} + \frac{1}{2} m \omega_n^2 (q_j - q_{j+1})^2 + V(q_j) \right]$$

and $\beta_n = \beta/n$ and $\omega_n = 1/(\beta_n \hbar)$ gives

$$\langle E \rangle = \frac{1}{(2\pi\hbar)^n Q} \int d\mathbf{p} \int d\mathbf{q} e^{-\beta_n H(\mathbf{p}, \mathbf{q})} \mathcal{E}(\mathbf{q}) \equiv \langle \mathcal{E}(\mathbf{q}) \rangle,$$

where

$$\mathcal{E}(\mathbf{q}) = \frac{1}{2\beta_n} - \frac{1}{2n} \sum_{j=1}^n m \omega_n^2 (q_j - q_{j+1})^2 + \frac{1}{n} \sum_{j=1}^n V(q_j) \equiv \mathcal{T}(\mathbf{q}) + \mathcal{V}(\mathbf{q}).$$

Centroid virial estimator

So the thermodynamic energy estimator

$$\mathcal{E}_{\text{TD}}(\mathbf{q}) = \frac{1}{2\beta_n} - \frac{1}{2n} \sum_{j=1}^n m\omega_n^2 (q_j - q_{j+1})^2 + \frac{1}{n} \sum_{j=1}^n V(q_j)$$

can be derived in two different ways, both of which show that its ring polymer average $\langle \mathcal{E}_{\text{TD}}(\mathbf{q}) \rangle$ will give the correct result $\langle E \rangle = \text{tr} [e^{-\beta \hat{H}} \hat{H}] / Q$ for the average energy in the canonical ensemble.


But this is not the only estimator that will do so. The *centroid virial* energy estimator^{5,6}

$$\mathcal{E}_{\text{CV}}(\mathbf{q}) = \frac{1}{2\beta} + \frac{1}{2n} \sum_{j=1}^n (q_j - \bar{q}) \frac{dV(q_j)}{dq_j} + \frac{1}{n} \sum_{j=1}^n V(q_j),$$


where $\bar{q} = (1/n) \sum_{j=1}^n q_j$, can be shown to have the same average $\langle \mathcal{E}_{\text{CV}}(\mathbf{q}) \rangle = \langle \mathcal{E}_{\text{TD}}(\mathbf{q}) \rangle$ as the thermodynamic energy estimator, with a *way* smaller variance $\Delta E = \sigma_E^2 = \langle \mathcal{E}(\mathbf{q})^2 \rangle - \langle \mathcal{E}(\mathbf{q}) \rangle^2$:

Comparison:

Thermodynamic Energy Estimator
(SHO with $h\nu/kT = 10$)



Centroid Virial Estimator
(SHO with $h\nu/kT = 10$)



Note that:

1. $\langle E \rangle$ is the same in both cases, and converges on the correct result with an error of $O(n^{-2})$.
2. $\langle E \rangle$ is converged to graphical accuracy by the time $n \simeq 5h\nu/kT = 5\beta\hbar\omega$.
3. The standard deviation of the thermodynamic estimator increases asymptotically as $n^{1/2}$.
4. If $\langle E \rangle$ were calculated using this estimator by Monte Carlo integration, the required number of samples M would increase linearly with n (because the standard error in the mean is proportional to $\sigma_E/M^{1/2}$).
5. By contrast, the standard deviation of the centroid virial estimator is *independent of n* for large n .

3. Integrating the equations of motion⁷

The standard way to integrate classical trajectories in molecular dynamics

$$\dot{\mathbf{p}} = -\frac{\partial H(\mathbf{p}, \mathbf{q})}{\partial \mathbf{q}} = -\frac{\partial V(\mathbf{q})}{\partial \mathbf{q}}$$

$$\dot{\mathbf{q}} = +\frac{\partial H(\mathbf{p}, \mathbf{q})}{\partial \mathbf{p}} = +\frac{\mathbf{p}}{m}$$

is to use the velocity Verlet algorithm ($e^{-\mathcal{L}\delta t} \simeq e^{-\mathcal{L}_V\delta t/2}e^{-\mathcal{L}_T\delta t}e^{-\mathcal{L}_V\delta t/2}$):

$$\mathbf{p} := \mathbf{p} - \frac{\delta t}{2} \frac{\partial V(\mathbf{q})}{\partial \mathbf{q}}, \quad \text{exact evolution under } H = V \text{ for time } \delta t/2$$

$$\mathbf{q} := \mathbf{q} + \delta t \frac{\mathbf{p}}{m}, \quad \text{exact evolution under } H = T \text{ for time } \delta t$$

$$\mathbf{p} := \mathbf{p} - \frac{\delta t}{2} \frac{\partial V(\mathbf{q})}{\partial \mathbf{q}}, \quad \text{exact evolution under } H = V \text{ for time } \delta t/2.$$

For PIMD, one could write the ring polymer Hamiltonian as

$$H_n(\mathbf{p}, \mathbf{q}) = \frac{|\mathbf{p}|^2}{2m} + V(\mathbf{q})$$

with

$$V(\mathbf{q}) = \sum_{j=1}^n \left[\frac{1}{2} m \omega_n^2 (q_j - q_{j+1})^2 + V(q_j) \right]$$

and use the standard velocity Verlet algorithm.

However, this would require a very small time step δt because of the stiff harmonic springs between the beads [$\omega_n = n/(\beta\hbar)$].

So we prefer to write

$$H_n(\mathbf{p}, \mathbf{q}) = H_0(\mathbf{p}, \mathbf{q}) + V(\mathbf{q})$$

where

$$H_0(\mathbf{p}, \mathbf{q}) = \sum_{j=1}^n \left[\frac{p_j^2}{2m} + \frac{1}{2} m \omega_n^2 (q_j - q_{j+1})^2 \right]$$

and

$$V(\mathbf{q}) = \sum_{j=1}^n V(q_j),$$

and to use the following time evolution algorithm:

$$e^{-\mathcal{L}\delta t} \simeq e^{-\mathcal{L}_V \delta t / 2} e^{-\mathcal{L}_0 \delta t} e^{-\mathcal{L}_V \delta t / 2}.$$

Because it is harmonic,

$$H_0(\mathbf{p}, \mathbf{q}) = \sum_{j=1}^n \left[\frac{p_j^2}{2m} + \frac{1}{2} m \omega_n^2 (q_j - q_{j+1})^2 \right]$$

can be diagonalised with a normal mode transformation

$$\tilde{p}_k = \sum_{j=1}^n p_j C_{jk} \quad \text{and} \quad \tilde{q}_k = \sum_{j=1}^n q_j C_{jk}$$

$$C_{jk} = \begin{cases} \sqrt{1/n}, & k = 0 \\ \sqrt{2/n} \cos(2\pi jk/n), & k = 1 \dots n/2 - 1 \\ \sqrt{1/n} (-1)^j, & k = n/2 \\ \sqrt{2/n} \sin(2\pi jk/n), & k = n/2 + 1 \dots n - 1, \end{cases}$$

which gives

$$H_0(\tilde{\mathbf{p}}, \tilde{\mathbf{q}}) = \sum_{k=0}^{n-1} \left[\frac{\tilde{p}_k^2}{2m} + \frac{1}{2} m \omega_k^2 \tilde{q}_k^2 \right]$$

with


$$\omega_k = 2\omega_n \sin(k\pi/n).$$

Normal mode transformation

In fact

$$\sum_{j=1}^n (q_j - q_{j+1})^2 = \mathbf{q}^T (2\mathbf{I} - \mathbf{A})\mathbf{q},$$

where \mathbf{A} is the adjacency matrix of the cyclic hydrocarbon C_nH_n in Hückel theory. So the ring polymer normal mode transformation corresponds to doing a simple HMO calculation:



Moreover

$$\tilde{p}_k = \sum_{j=1}^n p_j C_{jk} \quad \text{and} \quad \tilde{q}_k = \sum_{j=1}^n q_j C_{jk}$$

is just a pair of discrete Fourier transforms, which can be done very efficiently for large n using the FFT algorithm.

Ring polymer evolution

Bringing all of this together, the (microcanonical) ring polymer evolution

$$e^{-\mathcal{L}\delta t} = e^{-\mathcal{L}_V\delta t/2} e^{-\mathcal{L}_0\delta t} e^{-\mathcal{L}_V\delta t/2}$$

proceeds as follows:⁷

$$p_j := p_j - \frac{\delta t}{2} \frac{dV(q_j)}{dq_j}$$

$$\tilde{p}_k := \sum_{j=1}^n p_j C_{jk} \quad \tilde{q}_k := \sum_{j=1}^n q_j C_{jk}$$

$$\begin{pmatrix} \tilde{p}_k \\ \tilde{q}_k \end{pmatrix} := \begin{pmatrix} \cos \omega_k \delta t & -m\omega_k \sin \omega_k \delta t \\ (1/m\omega_k) \sin \omega_k \delta t & \cos \omega_k \delta t \end{pmatrix} \begin{pmatrix} p_k \\ q_k \end{pmatrix}$$

$$p_j := \sum_{k=0}^{n-1} C_{jk} \tilde{p}_k \quad q_j := \sum_{k=0}^{n-1} C_{jk} \tilde{q}_k$$

$$p_j := p_j - \frac{\delta t}{2} \frac{dV(q_j)}{dq_j}.$$

Non-ergodicity

These microcanonical ring polymer trajectories are all very well, but they are no good (on their own) for calculating thermal averages such as

$$\langle V \rangle = \frac{1}{(2\pi\hbar)^n Q} \int d\mathbf{p} \int d\mathbf{q} e^{-\beta_n H_n(\mathbf{p}, \mathbf{q})} \mathcal{V}(\mathbf{q}),$$

for two reasons:

1. They conserve $H_n(\mathbf{p}, \mathbf{q})$. So they do not explore all (Boltzmann-weighted) values of $H_n(\mathbf{p}, \mathbf{q})$.
2. They are far from ergodic.⁸ (E.g., for the SHO potential, $V(q) = \frac{1}{2}m\omega^2 q^2$, $H_n(\mathbf{p}, \mathbf{q})$ is diagonal in the normal mode representation; there is no energy flow between the normal modes. And for a mildly anharmonic potential, one would have to run a microcanonical trajectory for an awfully long time to see any energy flow.)

The PILE thermostat

Both problems can be fixed by attaching a “path integral Langevin equation” (PILE) thermostat to the dynamics. I.e., by replacing

$$e^{-\mathcal{L}_V \delta t/2} e^{-\mathcal{L}_0 \delta t} e^{-\mathcal{L}_V \delta t/2}$$

with

$$e^{-\mathcal{L}_\gamma \delta t/2} e^{-\mathcal{L}_V \delta t/2} e^{-\mathcal{L}_0 \delta t} e^{-\mathcal{L}_V \delta t/2} e^{-\mathcal{L}_\gamma \delta t/2},$$

in which the thermostating ($e^{-\mathcal{L}_\gamma \delta t/2}$) steps are implemented as follows:⁷

$$\tilde{p}_k := \sum_{j=1}^n p_j C_{jk}$$

$$\tilde{p}_k := e^{-\gamma_k \delta t/2} \tilde{p}_k + \sqrt{m(1 - e^{-\gamma_k \delta t})/\beta_n} \xi_k$$

$$p_j = \sum_{k=0}^{n-1} C_{jk} \tilde{p}_k.$$

Here ξ_k is an independent Gaussian number (a normal deviate with zero mean and unit variance) that is different for each invocation of $e^{-\mathcal{L}_\gamma \delta t/2}$.

The PILE algorithm corresponds to attaching a separate Langevin thermostat to each internal mode of the free ring polymer,

$$\frac{d}{dt}\tilde{q}_k = \frac{\tilde{p}_k}{m}$$

$$\frac{d}{dt}\tilde{p}_k = -m\omega_k^2\tilde{q}_k - \gamma_k\tilde{p}_k + \sqrt{\frac{2m\gamma_k}{\beta_n}}\xi_k(t),$$

where $\xi_k(t)$ represents an uncorrelated, Gaussian-distributed random force with unit variance and zero mean [$\langle\xi_k(t)\rangle = 0$ and $\langle\xi_k(0)\xi_k(t)\rangle = \delta(t)$].

The autocorrelation time

$$\tau_V = \frac{1}{\langle V^2 \rangle - \langle V \rangle^2} \int_0^\infty \langle (V(0) - \langle V \rangle)(V(t) - \langle V \rangle) \rangle dt$$

of the free ring polymer normal mode potential $V = m\omega_k^2 \tilde{q}_k^2/2$ can be worked out analytically for this Langevin dynamics and is⁹

$$\tau_V = \frac{1}{2\gamma_k} + \frac{\gamma_k}{2\omega_k^2}$$

for $k > 0$ [$\omega_k > 0$]. The optimum friction coefficient γ_k (which minimises τ_V) is therefore simply $\gamma_k = \omega_k$ for $k > 0$, leaving a single physical parameter τ_0 to be specified for thermostating the centroid mode ($k = 0$):

$$\gamma_k = \begin{cases} 1/\tau_0, & k = 0, \\ \omega_k, & k > 0. \end{cases}$$

4. Multidimensional generalisation

The above equations have been given for a simple one-dimensional problem with

$$\hat{H} = \frac{\hat{p}^2}{2m} + V(\hat{q}).$$


However, in the absence of identical particle (fermionic and bosonic) exchange effects, they are straightforward to generalise to a multidimensional Hamiltonian of the form

$$\hat{H} = \sum_{i=1}^N \frac{|\hat{\mathbf{p}}_i|^2}{2m_i} + V(\hat{\mathbf{r}}_1, \dots, \hat{\mathbf{r}}_N).$$

For example, the ring polymer Hamiltonian becomes

$$H_n(\{\mathbf{p}\}, \{\mathbf{r}\}) = \sum_{i=1}^N \sum_{j=1}^n \left(\frac{|\mathbf{p}_{ij}|^2}{2m_i} + \frac{1}{2} m_i \omega_n^2 |\mathbf{r}_{ij} - \mathbf{r}_{ij+1}|^2 \right) + \sum_{j=1}^n V(\mathbf{r}_{1j}, \dots, \mathbf{r}_{Nj}).$$

E.g., for $N=2$ particles and $n=5$ beads:



————— physical potential interactions

- - - - - harmonic springs between beads

Identical particle exchange effects

Identical particle exchange effects become important when the de Broglie thermal wavelengths $\Lambda_i(T) = h/\sqrt{2\pi m_i kT}$ exceed the hard sphere diameters of the atoms.

These effects can in principle be included by considering dimerisation, trimerisation, etc. of ring polymers (see Chandler and Wolynes²).

However, it is hardly ever necessary for those of us who work in chemistry departments to have to worry about them, because (e.g.) these effects are negligible in liquid para-hydrogen even at its melting temperature (13.8 K).

5. References

1. R. P. Feynman and A. R. Hibbs, *Quantum Mechanics and Path Integrals* (McGraw-Hill, New York, 1965).
2. D. Chandler and P. G. Wolynes, *J. Chem. Phys.* **74**, 4078 (1981).
3. P. A. M. Dirac, *The Principles of Quantum Mechanics* (4th Edition, Oxford University Press, Oxford, 1958).
4. M. Parrinello and A. Rahman, *J. Chem. Phys.* **80**, 860 (1984).
5. M. F. Herman, E. J. Bruskin and B. J. Berne, *J. Chem. Phys.* **76**, 5150 (1982).
6. The centroid virial energy estimator is derived in the present notation, and extended to give an estimator for the imaginary time velocity auto-correlation function, in Appendix A of S. Habershon, B. J. Braams and D. E. Manolopoulos, *J. Chem. Phys.* **127**, 174108 (2007).
7. M. Ceriotti, M. Parrinello, T. E. Markland and D. E. Manolopoulos, *J. Chem. Phys.* **133**, 124104 (2010).
8. R. W. Hall and B. J. Berne, *J. Chem. Phys.* **81**, 3641 (1984).
9. M. Ceriotti, Ph.D. thesis, ETH Zurich (2010).